

## **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.



# RTD UPDATES: Farm Chemical Use

## Data updates from the Resources and Technology Division

Economic Research Service  
U.S. Department of Agriculture

March 1994  
Number 2

### Chemical Use on Oranges and Water Quality Risk

- Pest scouting to determine the need for pesticide treatments, a fundamental element of Integrated Pest Management programs, occurred on most surveyed orange acreage in the three states.
- Soil nutrient testing took place on 70 percent or more of surveyed orange acreage in Florida; in California/Arizona, leaf testing for nutrients was relatively more common.
- The most widespread potential risk to water quality from surveyed farms comes from applied chemicals which may dissolve in runoff and be carried to nearby lakes and streams.

This issue of RTD Updates summarizes findings on orange production practices related to pest control and nutrient management from the 1991 Fruit Chemical Use Survey. It reports field level practices for a sample of approximately 300 orange growers in Arizona, California and Florida, and links these practices to soils characteristics to assess the potential risks to water quality. The Fruit Chemical Use Survey is part of the USDA Pesticide Data Program, a cooperative effort between several USDA agencies, EPA and FDA to produce statistically reliable information on pesticide use.

Scouting and nutrient management, key practices which can reduce the risks of agricultural chemical use to water quality and human and environmental health, are already

adopted on a majority of surveyed orange acreage. Other alternatives, such as the release of beneficial insects, are less widely adopted. The survey shows potential for further adoption of other chemical-reducing practices.

Pesticide applicator labor hours are a measure of the potential human health risks which can arise from direct exposure to pesticides on the farm. Among surveyed farms in California/Arizona, about half of the total hours worked by pesticide applicators were on farms using high amounts of Class I (most toxic) pesticides, compared to 28 percent on farms using no Class I pesticides.


Combined chemical and soil properties determine the risks to water quality from chemical use. Survey results indicate little potential risk for chemicals applied on orange acreage to leach into groundwater, with only 7 percent of surveyed farms falling into the highest risk category. Surface water is at higher risk, with 23 and 44 percent, respectively, of surveyed farms in the highest risk categories for chemicals to attach to eroding soil particles and dissolve in runoff.

Contacts: Steve Crutchfield (202) 219-0444  
Ann Vandeman (202) 219-0434

#### About RTD UPDATES

RTD UPDATES is a semimonthly series featuring data relating to agricultural resources, the environment, food safety, and technology. These UPDATES report recent data from surveys of farm operators and others knowledgeable about changing agricultural resource conditions, with only minimal interpretation or analysis. Please contact the individual listed at the end of the text for additional information about the data in this UPDATE. If you would like to be added to the mailing list or have other questions about RTD UPDATES, contact Richard Magleby, (202) 219-0436.

Scouting use by surveyed orange growers





**Table 1--Pesticide active ingredients applied to surveyed orange acreage, by toxicity class, 1991\***

Class I pesticides	Class II pesticides	Class III pesticides
2,4-D	Abamectin	Benomyl
Aldicarb	Basic copper sulfate	Bromacil
Azinphos-methyl	Captafol	Copper ammonium complex
Chloropicrin	Carbaryl	Cryolite
Copper hydroxide	Chlorophacinone	EPTC
Copper sulfate	Chlorpyrifos	Ferbam
Cyfluthrin	Diazinon	Gibberellic acid
Diphacinone	Dicofol	Malathion
Fenamiphos	Dimethoate	Metalddehyde
Fenbutatin-oxide	Diquat	Metam-sodium
Fluazifop-P-butyl	Diuron	MSMA
Formetanate hydrochloride	Ethion	NAA
Fosetyl-al	Fenpropathrin	Napropamide
Metalaxyl	Fluvalinate	Norflurazon
Methidathion	Glyphosate	Oryzalin
Methomyl	Methiocarb	Oxydemeton-methyl
Methyl bromide	Oxyflourfen	Piperonyl butoxide
Mevinphos	Oxythioquinox	Pyrethrins
Naled	Pendimethalin	Ryania
Paraquat	Phosmet	Sabadilla
Propargite		Sethoxydim
Strychnine		Simazine
		Sulfur
		Trifluralin

Pesticide active ingredients are classified by EPA into four toxicity categories, according to their acute toxic affects, from most acutely toxic (Class I) to least toxic (Class IV). Source: Farm Chemicals Handbook '91, Mesiter Publishing Company, Willoughby, OH.

\* Three pesticide active ingredients used on oranges, Copper oxychloride sulfate, Phytophthora palmivora, and Potassium salts, are either exempt from residue tolerance or lacking mammalian toxicity. One pesticide, Petroleum Distillate, was ranked as a Class IV chemical.

**Table 2--Surveyed orange acreage and pesticide applicator labor hours, by use of Toxicity Class I pesticides, 1991**

State/crop	High use	Low use	No use	High use	Low use	No use
	Percent of surveyed acreage			Percent of applicator hours		
Florida - fresh	2	96	3	*	99	1
Florida - processing	12	64	23	12	78	11
AZ/CA - fresh	52	28	20	52	20	28

\* Less than 1 percent; high use = Class I pesticides make up more than 24 percent of total lbs. of a.i. applied; low use = Class I pesticides make up 1-24 percent of total lbs. of a.i. applied; no use = no Class I pesticides applied.

Table reads, e.g.: "Farms with high use of Class I pesticides account for 12 percent of all pesticide applicator hours on processing orange acreage in Florida."

**Table 3--Other pest management practices on surveyed orange acreage, 1991**

State/crop	Beneficial insects	Trap crops	Pheromones	Pruning/canopy mgmt.	Debris removal	Reduced pesticide use	Irrigation	Tillage/mowing	Pest resistant varieties
	Percent of surveyed acreage								
AZ/CA - fresh	19	8	62	50	59	45	17	28	16
Florida - fresh	58	0	2	29	66	88	6	84	12
Florida - processing	4	3	0	22	39	20	20	65	9

Table reads, e.g.: "19 percent of orange acreage on surveyed farms in California and Arizona was treated with beneficial insects."



**Table 4--Scouting use and pesticide application decision factors for oranges, 1991**

Item	Arizona/California-fresh		Florida-fresh		Florida-processing	
	Scouted	Not scouted	Scouted	Not scouted	Scouted	Not scouted
Percent of surveyed acreage						
Use of scouting	69	31	81	19	87	13
Application decision factor:	Percent of scouted/not scouted acreage					
Predetermined schedule	1	13	0	0	3	7
Scouting reports	90	1*	95	0	95	1*
Other	9	86	5	100	2	92

Table reads, e.g.: "69% of orange acreage on surveyed farms in California and Arizona was scouted. Scouting reports were the most important factor in deciding when to apply pesticides on 90% of this acreage."

\*Growers who did not report scouting their fields may rely on a neighbor who scouts or on local reports provided by extension.

**Table 5--Characteristics of surveyed orange growers by use of scouting, 1991**

Characteristic	Arizona/California-fresh		Florida-fresh		Florida-processing	
	Scouted	Not scouted	Scouted	Not scouted	Scouted	Not scouted
Percent of surveyed growers						
Some college education	83	61	73	83	82	67
Income over \$100,000	73	81	18	50	29	67
Certified restricted pesticide applicator	50	34	64	50	39	67
Average age of operator	54	52	47	56	52	57
Average years of experience	25	21	18	29	24	29

Table reads, e.g.: "73% of surveyed growers in California and Arizona who used scouting had gross farm sales of more than \$100,000."

Percent of fruit acres on surveyed orange farms to which pesticides or fertilizers are applied through irrigation water




Chart reads, e.g.: "Surveyed orange growers in Florida fertigated 15% of their fruit acreage."

Nutrient management practices used on surveyed orange acreage




Chart reads, e.g.: "Soil testing was performed on 27 percent of orange acreage on surveyed farms in California and Arizona"



**Distribution of potential water quality risk  
from pesticide use on surveyed orange acreage**




Chart reads, e.g., "More surveyed acreage was at highest risk for chemicals to dissolve in cropland runoff than for chemicals to attach to eroding soil particles." "Safe" or "Lowest" risk levels indicate a low or negligible probability of chemical contamination, while "Hazardous" or "Risky" cropland is most susceptible to water pollution from pesticide use on oranges.

Source: 1991 Fruit Chemical Use Survey, Economic Research Service and National Agricultural Statistics Service, USDA

**RTD UPDATES**

Economic Research Service  
U.S. Department of Agriculture  
1301 New York Avenue, N.W., Room 524  
Washington, DC 20005-4788

FIRST CLASS  
POSTAGE & FEES PAID  
USDA  
PERMIT NO. G-145

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-5881 (voice) or (202) 720-7808 (TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, DC 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.